



Enter Keyword(s)

All News

SEARCH

**Today's Medical & Research News**

Life Sciences

**New life sciences research from University of Alicante outlined**

June 10th, 2008

[Printer-Friendly Version](#)
[Email this Article](#)

[Click Here -- Agriculture -- Chen](#)
**MEDICAL NEWS**

[Professional](#)  
[Industry Executives](#)  
[Consumers](#)  
[Marketing/PR Professionals](#)

**NEWSRX LIBRARY**

[View the NewsRx Library](#)

**OUR PRODUCTS**

[Newsletters for Purchase](#)  
[NewsRx Passes](#)  
[NewsRx Bundles](#)  
[Custom Reports](#)  
[Site Licenses](#)

**MARKET RESEARCH**

[Business Intelligence Reports](#)  
[Competitive Intelligence Database](#)

[Privacy Policy](#)

**Marketing and PR Pros**

[Post press releases](#) [Learn more](#)

[PR Posting login](#)

(NewsRx.com) -- Fresh data on life sciences are presented in the report 'Whole cell electrochemistry of electricity-producing microorganisms evidence an adaptation for optimal exocellular electron transport.' "The mechanism(s) by which electricity-producing microorganisms interact with an electrode is poorly understood. Outer membrane cytochromes and conductive pili are being considered as possible players, but the available information does not concur to a consensus mechanism yet," scientists in Alicante, Spain report.

"In this work we demonstrate that *Geobacter sulfurreducens* cells are able to change the way in which they exchange electrons with an electrode as a response to changes in the applied electrode potential. After several hours of polarization at 0.1 V Ag/AgCl-KCl (saturated), the voltammetric signature of the attached cells showed a single redox pair with a formal redox potential of about -0.08 V as calculated from chronopotentiometric analysis. A similar signal was obtained from cells adapted to 0.4 V. However, new redox couples were detected after conditioning at 0.6 V. A large oxidation process beyond 0.5 V transferring a higher current than that obtained at 0.1 V was found to be associated with two reduction waves at 0.23 and 0.50 V. The apparent equilibrium potential of these new processes was estimated to be at about 0.48 V from programmed current potentiometric results. Importantly, when polarization was lowered again to 0.1 V for 18 additional hours, the signals obtained at 0.6 V were found to greatly diminish in amplitude, whereas those previously found at the lower conditioning potential were recovered," wrote J.P. Busalmen and colleagues, University of Alicante.

The researchers concluded: "Results clearly show the reversibility of cell adaptation to the electrode potential and point to the polarization potential as a key variable to optimize energy production from an electricity producing population."

Busalmen and colleagues published their study in *Environmental Science & Technology* (Whole cell electrochemistry of electricity-producing microorganisms evidence an adaptation for optimal exocellular electron transport. *Environmental Science & Technology*, 2008;42(7):2445-50).

For more information, contact J.P. Busalmen, Universidad de Alicante, Instituto de Electroquímica, Apartado de correos 99, 03080, Alicante, Spain.

Publisher contact information for the journal *Environmental Science & Technology* is: American Chemical Society, 1155 16th St., NW, Washington, DC 20036, USA.

Keywords: Spain, Alicante, Life Sciences.

This article was prepared by NewsRx editors from staff and other reports. Copyright 2008, NewsRx.com.

**NewsRx Article****RELATED TOPICS**

[Spain](#)  
[Alicante](#)  
[Life Sciences](#)

